



July 13, 2009

Hep C Doubles Risk for AIDS Illnesses

People coinfectd with both HIV and [hepatitis C virus \(HCV\)](#) have double the risk of developing an AIDS defining illness (ADI) as people infected with only HIV, according to a study [published](#) online July 10 in *Clinical Infectious Diseases*.

Numerous studies have demonstrated that HIV greatly accelerates HCV disease progression. It is a lot less clear, however, what impact HCV has on HIV disease progression. Studies have been mixed. Few studies have looked at the potential for an increased risk of developing an ADI in coinfectd people.

To determine the risk of ADIs in coinfectd individuals, Antonella d'Arminio, MD, at the University of Milan, and her colleagues examined the medical records of 5,397 HIV-positive patients in the ICONA Foundation Study Cohort. Most of the patients enrolled in the cohort in 1998, but some were enrolled as recently as 2008. Of those patients, 2,421 were also infected with HCV, and 2,976 were not. The average CD4 count was 418 in the coinfectd group and 458 in the group without HCV infection. Roughly one third were women.

There were 496 ADIs in the two groups. The rate of developing an ADI was two-fold higher in the coinfectd group, after adjusting for other risk factors. The rate of ADIs was three times as high for developing fungal infections, [wasting](#), [dementia](#) and bacterial infections such as [tuberculosis](#) and [Mycobacterium avium complex \(MAC\)](#). People taking antiretroviral (ARV) therapy were less likely to develop an ADI than people not taking ARV therapy. People with more severely damaged livers (cirrhosis) were also more likely to have an ADI.

The authors conclude that coinfectd people should be more carefully monitored for AIDS defining illnesses.